



FastPure®
Blood/Cell/Tissue/Bacteria
DNA Isolation Mini Kit

NB-54-0205

FastPure® Blood/Cell/Tissue/Bacteria DNA Isolation Mini Kit

Cat# NB-54-0205-01, NB-54-0205-02

Product Description

This kit is suitable for extracting genomic DNA from $\leq 200 \mu\text{l}$ of fresh or frozen anticoagulated whole blood, $< 25 \text{ mg}$ of animal tissue, $< 5 \times 10^6$ of cultured cells and $< 3 \times 10^9$ cultured bacterial samples. The kit is based on silica gel column purification technology that eliminates the need for extraction using phenol/chloroform organic solvents or time-consuming alcohol precipitation. With this kit, RNA, proteins, lipids and other inhibitory impurities can be removed at the greatest extent. The DNA obtained can be directly used in PCR, qPCR, restriction enzyme digestion and virus detection..

Components

Components	NB-54-0205-01 (50 rxns)	NB-54-0205-02 (200 rxns)
Buffer ACL	12 ml	48 ml
Proteinase K	1 ml	4 ml
Buffer BCL	12 ml	48 ml
Buffer WA	15 ml	60 ml
Buffer WB	20 ml	2 × 40 ml
Elution Buffer	20 ml	80 ml
FastPure gDNA Mini Columns II	50	2 × 100
Collection Tubes 2 ml	50	2 × 100

Buffer ACL: Sample lysis; Proteinase K: Perform enzymatic lysis of samples; Buffer BCL: Perform lysis of samples and provide the reaction environment; Buffer WA: Remove residual protein contamination; Buffer WB: Remove salt ions; Elution Buffer: Elute the bound DNA; FastPure gDNA Mini Columns II : Genomic DNA adsorption column; Collection Tubes 2 ml: Filtrate collection tubes

Storage

Apart from Proteinase K, store the other components at room temperature ($15 \sim 25^\circ\text{C}$); Proteinase K can be stored at $15 \sim 25^\circ\text{C}$ for 3 months. Store at $2 \sim 8^\circ\text{C}$ for long-term storage and transport at room temperature.

Applications

Fresh or frozen anticoagulated whole blood without nuclei ($\leq 200 \mu\text{l}$); Fresh or frozen anticoagulated whole blood with nuclei ($5 - 20 \mu\text{l}$); Cultured cells ($< 5 \times 10^6$); Animal tissue ($< 25 \text{ mg}$); Bacteria ($< 3 \times 10^9$).

Self-prepared Materials

PBS, RNase A (optional), Lysozyme (for Gram-positive bacteria extraction), absolute ethanol, sterilized 1.5 ml centrifuge tube, water bath, etc

Notes

1. Please add a specified amount of absolute ethanol to Buffer WA and Buffer WB according to the label before use.

Reagent Name	NB-54-0205-01 WA	NB-54-0205-01 WB	NB-54-0205-02 WA	NB-54-0205-02 WB
Volume of absolute ethanol (ml)	20	80	80	160

2. Avoid repeated freezing and thawing of samples, as this can result in genomic DNA degradation and a reduction in extraction efficiency.
3. Check if there is any precipitation in Buffer ACL, Buffer BCL and Buffer WA before use. If precipitates have formed, they can be re-dissolved in a 37°C-water bath and mixed well before use.
4. All procedures must be carried out at room temperature (15 ~ 25 °C).

Mechanism & Workflow



Bind genome DNA: Transfer the mixture to an adsorption column and centrifuge at 12,000 rpm (13,400 × g) for 1 min;

Removal of impurities such as protein: Add 500 µl Buffer WA and centrifuge at 12,000 rpm (13,400 × g) for 1 min;


Removal of salt ions: Add 600 µl Buffer WB and centrifuge at 12,000 rpm (13,400 × g) for 1 min (twice);

Removal of ethanol: Centrifuge the empty column at 12,000 rpm (13,400 × g) for 2 min, then open the lid and air-dry at room temperature for 2 - 5 min.

Provide optimal binding environment : Add 150 µl absolute ethanol and mix by vortexing.



Sample processing



Blood samples : Make up to 200 µl with Buffer ACL, then sequentially add 20 µl Proteinase K and 200 µl Buffer BCL. Mix by vortexing and incubate at 70°C for 10 min in a water bath;

Cell samples : Sequentially add 200 µl PBS, 20 µl Proteinase K, and 200 µl Buffer BCL. Mix by vortexing and incubate at 56°C for 10 min in a water bath;

Tissue samples : Sequentially add 200 µl Buffer ACL and 20 µl Proteinase K, then mix by vortexing and incubate at 56°C in a water bath until complete enzymolysis; add 200 µl Buffer BCL and mix by vortexing



Bacterial samples : (Gram-negative bacteria) Sequentially add 200 µl Buffer ACL, 20 µl Proteinase K and 200 µl Buffer BCL. Mix by vortexing and incubate at 56°C for 10 min a water bath;

(Gram-positive bacteria) Add 180 µl Lysozyme and incubate at 37°C for 30 min in a water bath, then sequentially add 20 µl Proteinase K and 200 µl Buffer BCL. Mix by vortexing and incubate at 56°C for 10 min in a water bath.



Elution : Add 50 - 200 µl Elution Buffer, incubate at room temperature for 2 - 5 min, then centrifuge at 12,000 rpm (13,400 × g) for 1 min.

Experiment Process

1/ Sample Processing

◇ Blood samples

1. Place blood sample ($\leq 200 \mu\text{l}$) in a 1.5 ml centrifuge tube. Add Buffer ACL to make up to $200 \mu\text{l}$, then shake to mix.
 - ▲ For samples of anticoagulated whole blood with nucleated red blood cells taken from poultry, birds and amphibians, the sample volume should be 5 - $20 \mu\text{l}$.
 - ▲ If the volume of blood is $>200 \mu\text{l}$, please use #NB-54-0017 (applicable range: 0.1 - 1 ml).
2. Sequentially add $20 \mu\text{l}$ Proteinase K and $200 \mu\text{l}$ Buffer BCL and shake to mix well.
3. Incubate at 70°C for 10 min in a water bath, mix by inverting several times. Ensure that there are no granular precipitates.
4. Proceed to **2/ Column Purification**.

◇ Cell samples

1. The total number of cells must not exceed 5×10^6 . Collect the cells by centrifuging at $300 \times g$ for 5 min, discarding the supernatant.
 - ▲ Adherent cells: Cell scrapers or clean pipette tips can be used, or digest the cells with trypsin, and then centrifuge to collect cells.
2. Sequentially add $200 \mu\text{l}$ PBS and $20 \mu\text{l}$ of Proteinase K, and shake to mix.
 - ▲ (Optional) If RNA residues significantly affect subsequent experiments, add $4 \mu\text{l}$ RNase A (100 mg/ml) in Step 2. Mix by inverting, and place at room temperature for 2 - 5 min.
3. Add $200 \mu\text{l}$ of Buffer BCL and shake to mix well. Incubate at 56°C for 10 min in a water bath. Mix by inverting several times. There may be genomic agglomerates in the solution, which is normal.
4. Proceed to **2/ Column Purification**.

◇ Tissue samples

1. Place $<25 \text{ mg}$ of chopped or ground tissues (For spleen and kidney, the amount should be less than 10 mg) into a 1.5 ml centrifuge tube. Then sequentially add $200 \mu\text{l}$ Buffer ACL and $20 \mu\text{l}$ Proteinase K, and shake to mix.
 - ▲ (Optional) If the RNA residues significantly affect subsequent experiments, add $4 \mu\text{l}$ RNase A (100 mg/ml) in Step 2. Mix by inverting, and place at room temperature for 2 - 5 min.
 - ▲ Excessive amount of samples will result in a reduction in DNA yield and purity. Samples such as liver, spleen and kidney are rich in DNA, and the samples shall be less than 10 mg in weight. For tissues with low DNA contents such as muscle and skin, samples can be increased to $20 \text{ mg} - 50 \text{ mg}$ in weight while proportionally increasing the amount of Buffer ACL, Proteinase K, Buffer BCL and absolute ethanol.
2. Place in a water bath at 56°C to complete enzymatic lysis. Mix by inverting several times. Ensure that there are no granular precipitates.
 - ▲ Mix by inversion to promote the lysis process. Chop the tissues into pieces to shorten the digestion time. Digestion time is dependent on the sample type and homogenization results. In general, 0.5 - 3 h is required for tissue samples, 6 - 8 h or overnight digestion is required for mouse tails.
 - ▲ If the tissue sample is ground into powder using liquid nitrogen, it can be extracted directly using the simple operating procedure for tissue samples.
3. Add $200 \mu\text{l}$ Buffer BCL, and shake to mix.
4. Proceed to **2/ Column Purification**.

- Simple operating procedure for tissue samples (This procedure can be used for tissue samples that are sufficiently ground into powder using liquid nitrogen)

1. Place $<25 \text{ mg}$ ground tissue (For spleen and kidney, the amount should be less than 10 mg) in a 1.5 ml centrifuge tube. Then sequentially add $200 \mu\text{l}$ Buffer ACL and $20 \mu\text{l}$ Proteinase K, and vortex until there are no granular precipitates.
 - ▲ (Optional) If RNA residues significantly affect subsequent experiments, $4 \mu\text{l}$ RNase A (100 mg/ml) (Vazyme #DE111) can be added in Step 1. Mix by inverting and place at room temperature for 2 - 5 min.
2. Add $200 \mu\text{l}$ Buffer BCL, and shake to mix.
3. Proceed to **08-2 Column Purification**.

◇ Bacteria samples Gram-negative bacteria

1. Centrifuge 1 - 5 ml of bacterial culture solution ($<3.0 \times 10^9$ bacteria) at 10,000 rpm ($11,500 \times g$) for 1 min, discarding the culture solution.

▲ The bacteria number can be measured by spectrophotometer. When $OD_{600} = 1.0$, the bacteria number is about 1.5×10^9 /ml.

2. Sequentially add 200 μ l Buffer ACL, 20 μ l Proteinase K, and 200 μ l Buffer BCL, and shake to mix.

3. Incubate at 56°C for 10 min in a water bath. Mix by inverting several times. Ensure that there are no granular precipitates.

▲ (Optional) If RNA residues significantly affect subsequent experiments, 4 μ l RNase A (100 mg/ml) can be added after Step 3. Then mix by inverting and place at room temperature for 2 - 5 min.

4. Proceed to **2/ Column Purification**.

Gram-positive bacteria

1. Centrifuge 1 - 5 ml of bacterial culture solution ($<3.0 \times 10^9$ bacteria) at 10,000 rpm ($11,500 \times g$) for 1 min, discarding the culture solution.

▲ The bacteria number can be measured by spectrophotometer. When $OD_{600} = 1.0$, the bacteria number is about 1.5×10^9 /ml

2. Add 180 μ l Lysozyme (20 mg/ml), shake to re-suspend the bacteria, and incubate at 37°C for 30 min in a water bath.

▲ The cell walls of most bacteria will break down sufficiently after 30 min in the water bath, but certain bacteria with thicker cell walls (such as Staphylococcus aureus) need to be treated for 1 - 2 h to completely break down cell walls. Please adjust the incubation time according to the types of bacteria.

▲ Gram-positive cocci with particularly thick cell walls can be extracted using the complex bacteria extraction procedure.

3. Sequentially add 20 μ l Proteinase K and 200 μ l Buffer BCL, and shake to mix.

▲ (Optional) If the RNA residues significantly affect subsequent experiments, 4 μ l RNase A (100 mg/ml) can be added in Step 3, then mix by inverting and place at room temperature for 2 - 5 min.

4. Incubate at 56°C for 10 min in a water bath. Mix by inverting several times. Ensure that there are no granular precipitates.

5. Proceed to **2/ Column Purification**

● Complex bacteria (Gram-positive cocci with particularly thick cell walls)

1. Centrifuge 1 - 5 ml of bacterial culture solution ($<3.0 \times 10^9$ bacteria) at 10,000 rpm ($11,500 \times g$) for 1 min, discarding the culture solution.

▲ The bacteria number can be measured by spectrophotometer. When $OD_{600} = 1.0$, the bacteria number is about 1.5×10^9 /ml.

2. Add 1 ml 70% ethanol and shake to re-suspend the bacteria, then place in an ice bath for 20 min. Centrifuge at 10,000 rpm ($11,500 \times g$) for 1 min, discarding the supernatant.

3. Add 180 μ l Lysozyme, shake to re-suspend the bacteria, and incubate in a water bath at 37°C until the cell walls are broken down. Centrifuge at 10,000 rpm ($11,500 \times g$) for 1 min, discarding the supernatant.

▲ The cell walls of most bacteria will break down sufficiently after 30 min in the water bath, but certain Gram-positive cocci with particularly thick cell walls need to be treated for 3 h to completely break down their walls. Please adjust the incubation time according to the types of bacteria.

4. Sequentially add 200 μ l Buffer ACL and 20 μ l Proteinase K, shake to re-suspend the bacteria, and incubate at 56°C for 30 min in a water bath.

5. Add 200 μ l Buffer BCL, and shake to mix.

▲ (Optional) If RNA residues significantly affect subsequent experiments, add 4 μ l RNase A (100 mg/ml) in Step 2. Mix by inverting, and place at room temperature for 2 - 5 min.

6. Incubate at 70°C for 10 min in a water bath.

7. Proceed to **2/ Column Purification**.

2/ Column Purification

1. Add 150 μ l absolute ethanol, shake to mix. Flocculent precipitate may appear, then centrifuge briefly to collect the liquid.

2. Place the FastPure gDNA Mini Columns II in 2 ml collection tubes. Transfer the mixture above (including the precipitate) to the adsorption column. Centrifuge at 12,000 rpm ($13,400 \times g$) for 1 min.

3. Discard the flow-through and place the adsorption column in collection tubes. Add 500 μ l Buffer WA along the tube wall (Check whether absolute ethanol has been added) to the adsorption column, and centrifuge at 12,000 rpm ($13,400 \times g$) for 1 min.

4. Discard the flow-through and place the adsorption column in collection tubes. Add 600 μ l Buffer WB along the tube wall (Check whether absolute ethanol has been added), and centrifuge at 12,000 rpm ($13,400 \times g$) for 1 min, discarding the flow-through.

5. Repeat **Step 4**.

6. Place the adsorption column in the collection tube. Centrifuge the empty column at 12,000 rpm (13,400 × g) for 2 min.

▲ After centrifugation, open the lid and air-dry for 2 - 5 min to completely remove the residual ethanol.

7. Transfer the adsorption column to a new 1.5 ml centrifuge tube (self-provided). Add 50 - 200 µl Elution Buffer to the center of the adsorption column membrane and incubate at room temperature for 2 - 5 min. Centrifuge at 12,000 rpm (13,400 × g) for 1 min.

Note: The following steps can help to increase DNA yield

▲ Pre-heat the Elution Buffer to 55 °C before elution;

▲ The solution from the first elution can be re-added to the adsorption column when carrying out the elution.

8. Discard the adsorption column and store the DNA products at -20 °C. For long-term storage, please store at -70 °C to prevent degradation.

9/ FAQ & Troubleshooting

Common Issues	Reason	Solution
Clogged adsorption column	1.Excessive amount of sample	Please input samples according to the compatible range.
	2.Decrease in Proteinase K activity	Replace with new Proteinase K, which must be stored at 2°C to 8°C after use.
	3.Insufficient sample lysis	Increase the incubation time in a 56°C water bath and mix by inverting more frequently.
Low DNA yield	1.Samples were frozen and thawed more than once	Avoid repeated freezing and thawing of samples. It is recommended to use fresh samples or samples that have been thawed only once.
	2.Insufficient lysis of animal tissue	The tissue should be chopped into pieces or ground with liquid nitrogen; Mix the sample thoroughly with Buffer ACL and Proteinase K; Appropriately increase the incubation time in the 56°C water bath.
	3.Incomplete breakdown of Gram-positive bacteria cell walls	The amount of Lysozyme can be increased appropriately or the digestion time can be increased; Carry out the experiment according to complex bacteria operation procedure.
	4.Decrease in Proteinase K activity	Replace with new Proteinase K, which must be stored at 2°C to 8°C after use.
	5.Lysis mixture is not fully transferred to the adsorption column	After addition of absolute ethanol, flocculent precipitate will appear in the solution, which must be transferred to the adsorption column together with the solution.
	6.Elution issues	Please use the Elution Buffer provided in the kit. If ddH ₂ O or other eluents are used, ensure that the pH of the eluent is between 8.0 and 9.0. The Elution Buffer can be pre-heated to 55°C before elution to help improve the DNA yield.
	7.Low elution efficiency	The eluent must be added to the center of the membrane; increase the elution volume or the number of elution times.
	8.Absolute ethanol was not added to Buffer WA/WB	Add the specified volume of absolute ethanol to Buffer WA and Buffer WB as indicated in the label.
Low DNA purity	1.Protein contamination	Buffer WA was not used for rinsing, or the correct volume of absolute ethanol was not added to Buffer WA. Add the specified volume of absolute ethanol as indicated in the label.
	2.Contamination with ions	Buffer WB was not used for rinsing, or rinsing was only carried out once. Please use Buffer WB to rinse twice as per the instructions.
	3.RNA residue	RNase A was not added in the specified step, or the RNase A incubation time was too short. Please add RNase A in the step specified in the instructions. For samples with high RNA content, the incubation time can be increased appropriately.
	4.Ethanol residue	Before elution, the empty tube was not centrifuged. Please open the lid and air-dry for 2 - 5 min to completely remove the ethanol after centrifugation.